



ELECTROLYTES
GEL

Concentrate of **mineral salts** which allow **rapid reconstitution of electrolyte stocks** lost through sweating.



INDICATIONS

- Moderate, intense or prolonged muscular effort resulting in sweating.
- Long-term transport.



PACKAGING



60 ml
Available in pack of 3 syringes



DIRECTIONS FOR USE

Make fresh water available at all times.

ADULT HORSE (500 KG)	RECOMMENDED DAILY DOSE	WHEN TO ADMINISTER
Moderate effort, transport over long distances	½ to 1 syringe, or 6 to 12 ml/100 kg body weight	After the event
Short-term intense effort (racing)	1 to 2 syringes, or 12 to 24 ml/100 kg body weight	1 syringe in the evening after exercise. Repeat the next morning in case of heavy sweating.
Prolonged intense effort (endurance)	1 to 2 syringes, or 12 to 24 ml/100 kg body weight	1 syringe morning and/or evening up to 10 days following the event

For ponies, feed a daily dose corresponding to the animal's body weight.



Gel

* Free from Naturally
Occurring Prohibited
Substances (NOPS),
in accordance with
the regulations of the
racing codes, FEI, FFE
and SHF.



DETAILED COMPOSITION

Demineralized water, sodium chloride, dextrose, potassium chloride, fructose, sodium citrate, magnesium chloride, glycine, orange flavouring.

ANALYTICAL CONSTITUENTS

Humidity	45%
Total protein.....	1.5%
Crude ash.....	32%
Sodium	9%

1 SYRINGE OF ELECTROLYTES GEL (60 ml) PROVIDES



13,175 mg chloride, 6,975 mg sodium, 3,480 mg potassium and 275 mg magnesium.



CONSERVATION

- Store in a dry place, protected from light, at a temperature between 5 and 25°C.
- Shelf life: 24 months from the date of manufacture.



PROPERTIES

Ingestion of the **mineral salts** contained in **ELECTROLYTES GEL** allows rapid reconstitution of electrolyte stocks lost through sweating.

A 60 ml syringe can replace the chloride, sodium, potassium and magnesium losses of approximately 2.5 L of sweat.

Dextrose and **glycine** facilitate the absorption of sodium.



PRECAUTIONS FOR USE

Due to the high concentration of mineral salts in **Reverdy ELECTROLYTES GEL**, it is not recommended to distribute it on the day of an endurance race.

In fact, if the horse does not drink enough during the event, the ingestion of **Reverdy ELECTROLYTES GEL** is likely to aggravate the dehydration of the body by creating a call for water in the lumen of the digestive tract.

On the other hand, it is possible to distribute **REVERDY ELECTROLYTES LIQUID** or **POWDER** on the day of the race. They must be diluted in a large volume of water and made available in a container next to the usual bucket of fresh water.