

Locomotor system



FLEXY LIQUID

DESCRIPTION

A preparation based on the three major chondroprotective agents (chondroitin, glucosamine and MSM) and hyaluronic acid.

Contributes to relieving articular pains, slowing the evolution of arthritis and improving articular lubrication and cartilage elasticity.

INDICATIONS

Prevent articular problems.

REVERDY FLEXY LIQUID can be used at higher doses in the following situations:

- Sudden/considerable articular stress:
 - › breaking/pre-training;
 - › intensive training period;
 - › competition;
 - › work on hard ground.
- Articular pain/lameness due to arthritis in the aged horse;
- In compliment to articular surgery.



1 L

5 L

DIRECTIONS FOR USE

Distribute by mixing with cereals or the pelleted feed, or to be directly swallowed by administering into the mouth using a syringe. Shake well before use.

ADULT HORSE (500 KG)	DAILY DOSE
Loading dose (1 st month)	50 ml equivalent to 10 ml /100 kg (liveweight)
Maintenance dose (or preventive)	25 ml equivalent to 5 ml /100 kg (liveweight)
Maximum recommended dose	75 ml equivalent to 15 ml /100 kg (liveweight)

* Non doping product without Prohibited Natural Feed Substances (analysed within the framework of a control scheme).



Liquid

For ponies and youngstock, administer a daily dose in proportion to the animal's liveweight.

COMPOSITION

Demineralised water, glucosamine sulphate 2 KCl (marine origin), MSM (Methylsulfonylmethane), fructose, chondroitin sulphate (marine origin), hyaluronic acid, natural orange flavouring.

ANALYTICAL CONSTITUENTS

Humidity	75%
Total protein	5.5%
Ash	5.5%
Sodium	0.3%

50 ML OF FLEXY LIQUID PROVIDE

10,000 mg of glucosamine sulphate 2KCl, 4,000 mg of MSM, 2,000 mg of marine chondroitin sulphate and 300 mg of hyaluronic acid.

PROPERTIES

Chondroitin is a constituent of proteoglycans whose role are to maintain correct hydration of cartilage and bones. Furthermore, it directly protects cartilage cells from enzymatic reactions and free radicals.

Glucosamine is the precursor of many of the constituents of proteoglycans and of hyaluronic acid. In cartilage, hyaluronic acid is bound with proteoglycans and forms aggregates which assure good hydration of this tissue. In the synovial fluid of joints, hyaluronic acid has a role of lubricant and chondroprotective agent.


MSM also possesses chondroprotective properties. It is also a source of organic sulphur indispensable to the synthesis of collagen, an abundant cartilage protein, giving it hydration, resistance, elasticity and suppleness properties.

CONSERVATION

Store in a dry place, away from light, at room temperature.

Use within 30 days after opening.

Shelf life : 24 months.



Conceived by our research and development department.
Manufactured in our laboratory.